


RECONSTRUCTING THE RIGHT-HANDED NEUTRINO MASSES IN LEFT-RIGHT SYMMETRIC THEORIES

P. HOSTEINS ^a

*Service de Physique Théorique, CEA-Saclay
91191 Gif-sur-Yvette, France*



1 Introduction : reviewing the Seesaw Mechanism

A very popular mechanism to generate masses for the left-handed neutrinos is the so-called seesaw mechanism, which consists of coupling the neutrinos to heavy particles and integrating them out.

The most studied type of seesaw is the type I : we introduce 3 right handed neutrinos N_R (singlets of $SU(2)_L$) with a heavy Majorana mass matrix M_R and a usual Dirac coupling to the ν_L and the Higgs. When the Higgs gets a vev v we generate a mass for the light neutrinos :

$$m_\nu = -v^2 Y_\nu^T M_R^{-1} Y_\nu \quad (1)$$

There is also a type II seesaw, which introduces a scalar Δ_L triplet of $SU(2)_L$. We can write a coupling $\frac{1}{2} f_{Lij} l_{Li}^T \Delta_L C l_{Lj}$ with f_L symmetric. When the electrically neutral component of Δ_L gets a vev $\langle \Delta^0 \rangle = v_L$ we get a mass for the neutrinos :

$$m_\nu = v_L f_L \quad (2)$$

In some classes of models both types of contributions to the neutrino mass matrix are present. A frequent assumption is that one type of seesaw dominates over the other. We will consider

^ahosteins@cea.fr

that both types are present and solve for the heavy neutrino masses and mixings in this more general context. We then concentrate on the example of $SO(10)$.

2 Reconstruction procedure of the heavy neutrino mass spectrum

The general formula for the light neutrino masses we are considering is :

$$m_\nu = v_L f_L - v^2 Y_\nu^T M_R^{-1} Y_\nu \quad (3)$$

The theories which will allow us to solve for M_R are theories which contain a left-right (LR) symmetry. In this context there exists a gauge group $SU(2)_R$ which is a mirror of $SU(2)_L$ for the right-handed particles and the heavy Majorana mass derives from the same mechanism as the type II seesaw, with a triplet Δ_R of $SU(2)_R$. We have consequently $M_R = v_R f_R$, with $\langle \Delta^0 \rangle = v_R \gg v$, and the LR symmetry gives $f_L = f_R = f$ and Y_ν symmetric :

$$m_\nu = v_L f - \frac{v^2}{v_R} Y_\nu^T f^{-1} Y_\nu = \alpha f - \beta Y_\nu^T f^{-1} Y_\nu \quad (4)$$

Now to extract f we need to know Y_ν . Assuming Y_ν is known and we fix the neutrino low energy parameters, we can use $Z = Y_\nu^{-1/2} m_\nu (Y_\nu^{-1/2})^T$ and $X = Y_\nu^{-1/2} f (Y_\nu^{-1/2})^T$ to rewrite the above formula :

$$Z = \alpha X - \beta X^{-1} \quad Z \text{ and } X \text{ symmetric} \quad (5)$$

where Z is known. We can then use a mathematical trick to find the matrix X (hence f) as a function of Z , namely we diagonalise both sides with a complex orthogonal transformation $O_Z = O_X^b$. The above equation then translates to each of the eigenvalues :

$$x_i^\pm = \frac{z_i \pm \sqrt{4\alpha\beta + z_i^2}}{2\alpha} \quad (6)$$

There are 2 solutions for each eigenvalue, which means a total of 8 solutions for f , in agreement with¹. We label the solutions by the sign chosen for each x_i for example “+++”.

3 The example of $SO(10)$

We apply this procedure to supersymmetric $SO(10)$ theories with only symmetric mass matrices for the fermions and the following relations between the Yukawas : $Y_\nu = Y_u$ and $Y_e = Y_d$ (which means that we have two ten-dimensional representations in the Higgs sector).

3.1 Mass spectra

As explained earlier we find 8 different mass spectra and a richer phenomenology than the $SO(10)$ type I seesaw alone. Examples of such mass spectra are displayed in fig.1.

Our free parameters are v_R the B-L number breaking scale, at which the N_{Ri} masses are generated, the ratio β/α and many complex phases as we cannot rephase the fermions independently. We display the figures as functions of v_R and fix $\beta/\alpha = 1$. Although v_R is allowed to run from 10^{12}GeV to 10^{17}GeV we cut the values for which the $f_i > 1$ (perturbativity constraint). We also display dotted the region where the 2 contributions to m_ν begin to cancel each other with a fine-tuning smaller than 10%.

^bThis transformation is not a physical diagonalisation and is not always well defined, so one should use it with caution




Figure 1: Different mass spectra for the heavy neutrinos as a function of v_R (no complex phase) : the dotted part signals a tuning between the types I and II finer than 10%

4 solutions have a very small $M_1 \sim 10^5 \text{ GeV}$, just like in the pure type I $SO(10)$ seesaw, 2 have a quite stable $M_1 \sim 10^{9-10} \text{ GeV}$ and the last 2 have a quite heavy spectrum.

3.2 Leptogenesis

Having the masses and couplings of the heavy neutrinos we can also explore the results for leptogenesis and try to see if we can accomodate the matter-antimatter asymmetry of the Universe.

First we review the mechanism of leptogenesis : the N_R having complex couplings they do not decay symmetrically to leptons and antileptons. Considering that heavy neutrinos are created in the hot early universe they start decaying at $T \sim M_i$ creating a lepton asymmetry. The asymmetry created by the lightest is driven by the quantity^{3,4} :

$$\varepsilon_{\text{CP}} \simeq \frac{3}{8\pi} \frac{\Im[Y m_\nu^* Y^T]_{11}}{(Y Y^\dagger)_{11}} \frac{M_1}{v^2} \quad (7)$$

To find a good baryon asymmetry we must have $\varepsilon_{\text{CP}} > 10^{-6}$. We have computed ε_{CP} for the different spectra and display generic examples in fig.2.

We see that some solutions are unable to reproduce a correct asymmetry, in connection with the smallness of M_1 as ε_{CP} is proportionnal to it. Even taking into account flavour effects recently suggested^{6,7}, that could be relevant for models with hierarchical Y_ν , we are not able to accomodate sufficiently large values of ε_{CP} .

From the 4 remaining solutions, 2 of them can accomodate a good asymmetry for any value of the phases but they tend to favor heavy $M_1 (> 10^{10} \text{ GeV})$ which is in tension with constraints from gravitino production.




Figure 2: The 3 typical behaviours of the CP asymmetry (one majorana complex phase is $\pi/4$, the others are 0)

The last 2 give acceptable CP asymmetry for some specific values of the phases, including in the region with no tuning.

4 Conclusion

We have introduced a method to reconstruct the heavy neutrino mass matrix from the general typeI+typeII seesaw formula, postulating LR symmetry. In accordance with earlier work we found 8 different solutions, signing a richer phenomenology than pure type I or type II seesaw. Applying the procedure to $SO(10)$, we found that, in contrast with type I $SO(10)$ it is possible to accomodate a correct baryon asymmetry.

Finally, it is worth mentioning that the large regions of fine tuning in m_ν can be naturally explained if we extend $SO(10)$ to the bigger gauge group E_6 where the couplings Y_ν and f are exactly proportionnal at higher energies.

Acknowledgments

I would like to thank Stéphane Lavignac and Carlos Savoy for fruitful collaboration, as well as Michele Frigerio for useful advice. I am also grateful to the Moriond staff for allowing me to present this work and for the financial support.

References

1. E. K. Akhmedov and M. Frigerio, “Seesaw duality”, hep-ph/0509299
2. P. Hosteins, S. Lavignac, C. Savoy, “Quark-Lepton Unification and Eight-Fold Ambiguity in the Left-Right Symmetric Seesaw Mechanism”, hep-ph/0606078
3. T. Hambye and G. Senjanovic, “Consequences of triplet seesaw for leptogenesis”, Phys. Lett. B **582** (2004) 73 [hep-ph/0307237]
4. S. Antusch and S. F. King, “Type II leptogenesis and the neutrino massa scale”, Phys. Lett. B **597** (2004) 199 [hep-ph/0405093]
5. G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, “Towards a complete theory of thermal leptogenesis in the SM and MSSM”, Nucl. Phys. B **685** (2004) 89 [hep-ph/0310123]
6. O. Vives, “Flavoured leptogenesis: A successful thermal leptogenesis with $N(1)$ mass below $10^{8.5}$ -GeV”, Phys. Rev. D **73** (2006) 073006 [hep-ph/0512160]
7. P. Di Bari, “Seesaw geometry and leptogenesis”, Nucl. Phys. B **727** (2005) 318 [hep-ph/0502082]